

[Previous PDF](#)[Next PDF](#)

ELSEVIER

Journal of Chemical Neuroanatomy 75 (2016) 43–51

Contents lists available at ScienceDirect

Journal of Chemical Neuroanatomy

journal homepage: www.elsevier.com/locate/jchemneu

Review

Microwave frequency electromagnetic fields (EMFs) produce widespread neuropsychiatric effects including depression

Martin L. Pall

Professor Emeritus of Biochemistry and Basic Medical Sciences, Washington State University, 638 NE 41st Avenue, Portland, OR 97232-3312, USA



ARTICLE INFO

Article history:

Received 13 April 2015

Received in revised form 1 August 2015

Accepted 9 August 2015

Available online 21 August 2015

Keywords:

Excessive calcium effects

Oxidative/nitrosative stress

Low-intensity microwave electromagnetic fields

ABSTRACT

Non-thermal microwave/lower frequency electromagnetic fields (EMFs) act via voltage-gated calcium channel (VGCC) activation. Calcium channel blockers block EMF effects and several types of additional evidence confirm this mechanism. Low intensity microwave EMFs have been proposed to produce neuropsychiatric effects, sometimes called microwave syndrome, and the focus of this review is whether these are indeed well documented and consistent with the known mechanism(s) of action of such EMFs. VGCCs occur in very high densities throughout the nervous system and have near universal roles in release of neurotransmitters and neuroendocrine hormones. Soviet and Western literature shows that much of the impact of non-thermal microwave exposures in experimental animals occurs in the brain and peripheral nervous system, such that nervous system histology and function show diverse and substantial changes. These may be generated through roles of VGCC activation, producing excessive neurotransmitter/neuroendocrine release as well as oxidative/nitrosative stress and other responses. Excessive VGCC activity has been shown from genetic polymorphism studies to have roles in producing neuropsychiatric changes in humans. Two U.S. government reports from the 1970s to 1980s provide evidence for many neuropsychiatric effects of non-thermal microwave EMFs, based on occupational exposure studies. 18 more recent epidemiological studies, provide substantial evidence that microwave EMFs from cell/mobile phone base stations, excessive cell/mobile phone usage and from wireless smart meters can each produce similar patterns of neuropsychiatric effects, with several of these studies showing clear dose–response relationships. Lesser evidence from 6 additional studies suggests that short wave, radio station, occupational and digital TV antenna exposures may produce similar neuropsychiatric effects. Among the more commonly reported changes are sleep disturbance/insomnia, headache, depression/depressive symptoms, fatigue/tiredness, dysesthesia, concentration/attention dysfunction, memory changes, dizziness, irritability, loss of appetite/body weight, restlessness/anxiety, nausea, skin

Recommended Articles

Controversies related to electromagnetic field exposure on peripheral nerves

Ferhat Say, ... +7 ... , Ahmet Pişkin

Journal of Chemical Neuroanatomy • September 2016[Preview](#)[View PDF](#)[Save PDF](#)

Maternal exposure to a continuous 900-MHz electromagnetic field provokes neuronal loss and pathological changes in cerebellum of 32-day-old female rat offspring

Ersan Odacı, ... +7 ... , Orhan Baş

Journal of Chemical Neuroanatomy • September 2016[Preview](#)[View PDF](#)[Save PDF](#)

The 2100 MHz radiofrequency radiation of a 3G-mobile phone and the DNA oxidative damage in brain

Duygu Sahin, ... +5 ... , Nesrin Seyhan

Journal of Chemical Neuroanatomy • September 2016[Preview](#)[View PDF](#)[Save PDF](#)